

## 77. The Complex Role of Restoration in Wetland Greenhouse Gas Production

Susanna Theroux<sup>1\*</sup>([stheroux@lbl.gov](mailto:stheroux@lbl.gov)), Wyatt Hartman<sup>1</sup>, Susannah Tringe<sup>1</sup>

<sup>1</sup>DOE Joint Genome Institute, Walnut Creek, California

**Project Goals:** This project aims to identify the geochemical and biological controls on greenhouse gas (GHG) production in coastal wetlands. Current projections of wetland response to global climate change are poorly constrained due to a limited understanding of the microbial mediators of carbon cycling in wetland soils. Using large-scale environmental genome sequencing, we aim to identify the microbial sentinels of GHG production and consumption in historic and restored wetlands of the San Francisco Bay-Delta. An enhanced comprehension of microbial metabolic arsenals will allow us to recommend management practices for future wetland restoration projects to maximize carbon sequestration and minimize microbial production of greenhouse gases.

Wetland environments play a critical role in the global carbon cycle, storing up to 35% of all terrestrial carbon and producing up to 75% of all non-anthropogenic methane. However, estimates of wetland GHG budgets are difficult to constrain given large variations in salinity, tidal regimes, soil saturation, organic carbon content, restoration status, and other geochemical and biological variables. Belowground microbial communities are the main drivers of greenhouse gases (GHG) cycling and their response to climate change will dictate whether a wetland serves as a net carbon sink or source. Wetland restoration has been proposed as a potential long-term carbon sequestration strategy, however wetland site selection and management practices are critical for ensuring restored wetlands sequester more GHG than they emit.

In an effort to better understand the underlying factors that shape the balance of carbon flux in wetland soils, we targeted the microbial communities along a salinity gradient ranging from freshwater tidal marshes to hypersaline ponds in the San Francisco Bay-Delta region. Seawater delivers sulfate to wetland ecosystems, providing a natural barrier to methane production, and traditional dogma predicts a decrease in methane production as salt content increases. Using 16S rRNA gene sequencing and shotgun metagenomics, coupled with greenhouse gas monitoring, we sampled sixteen sites capturing a range of salinities and restoration statuses. Soils were characterized for a suite of geochemical parameters including soil moisture, percent organic carbon, pH, and micronutrient, cation, and metal concentrations. The results of our microbial diversity survey showed that sampling location, plant type, and salinity were the primary drivers of both methane production and belowground microbial community composition. As expected, freshwater wetland soils produced more methane than brackish and saline sites, also reflected in the increased relative abundance of methanogen populations. Notably, restored freshwater and brackish wetlands produced orders of magnitude more methane than their historic counterparts. And conversely, unrestored hypersaline ponds produced methane at rates rivaling those of brackish restored wetlands. Both 16S and metagenome sequencing allowed for the identification of hallmark methanogen species whose relative abundance trended consistently with methane production. Our study links belowground microbial communities with their aboveground greenhouse gas production and provides a benchmark for predicting wetland soil microbial response to the inevitable influence of climate change.

*The work conducted by the U.S. Department of Energy Joint Genome Institute is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. This project was funded by the DOE Early Career Research Program, grant number KP/CH57/1 to Susannah Green Tringe, and was also supported by DOE JGI Community Sequencing Program.*