

## Ectopic expression of *WRI1* affects fatty acid homeostasis in *Brachypodium distachyon* vegetative tissues

Yang Yang<sup>1,2,\*</sup>([young83@msu.edu](mailto:young83@msu.edu)), Jacob Munz<sup>3,4,†</sup>, Cynthia Cass<sup>3,4</sup>, Agnieszka Zienkiewicz<sup>1,2</sup>,  
Que Kong<sup>1,2</sup>, Wei Ma<sup>2,4</sup>, Sanjaya<sup>2,§</sup>, John Ohlrogge<sup>2,5</sup>, John Sedbrook<sup>3,4</sup>, **Christoph Benning**<sup>1,2,5</sup>

<sup>1</sup>Department of Biochemistry and Molecular Biology, Michigan State University, East Lansing, MI 48824, USA.

<sup>2</sup>Great Lakes Bioenergy Research Center, Michigan State University, East Lansing, MI 48824, USA.

<sup>3</sup> Great Lakes Bioenergy Research Center, Illinois State University, Normal, IL 61790, USA.

<sup>4</sup>School of Biological Sciences, Illinois State University, Normal, IL 61790, USA.

<sup>5</sup>Department of Plant Biology, Michigan State University, East Lansing, MI 48824, USA.

<sup>6</sup>MSU-Department of Energy, Plant Research Laboratory, East Lansing, MI 48824, USA.

<sup>§</sup>Present address: Energy and Environmental Science Institute, Department of Biology, West Virginia State University, Institute, WV 25112, USA.

<sup>†</sup>Present address: Department of Botany, University of British Columbia, Vancouver, BC V6T 1Z4, Canada.

<https://www.glbrc.org/research/plants>

**Project Goals:** This project aims to synthesize high-energy oils in vegetative tissues of biomass crops, including C4 grasses. Enhancing the energy content and liquid fuel yield of crop biomass can be achieved by accumulating triacylglycerols (TAG) in the crop. After extracting these compounds, lignocellulosic feedstock remains and can be used for processing and fermentation. Thus engineering oil production will result in novel biofuel crops that achieve multiple objectives. Our previous research has succeeded in producing TAG in *Arabidopsis* leaves and stems (Sanjaya et al., 2011; Sanjaya et al., 2013). Oil content was increased by combining transcription factors, down-regulation of starch synthesis, and addition of enzymes of TAG synthesis (Sanjaya et al., 2011). Testing these second generation approaches in the grass model plant *Brachypodium* is now efficient and cost-effective providing a ‘proof-of-concept’ strategy prior to moving into corn or other energy crops.

WRINKLED1 (*WRI1*) is a transcription factor, which governs fatty acid (FA) synthesis and indirectly TAG accumulation in oil storing plant tissues and ectopic expression of its cDNA has led to TAG accumulation in vegetative tissues of different dicotyledonous plants (Cernac and Benning, 2004; Cernac et al., 2006; Ma et al., 2013; Ma et al., 2015). The ectopic expression of *BdWRI1* in the grass *Brachypodium* (*Brachypodium distachyon*) induced the transcription of predicted genes involved in glycolysis and fatty acid (FA) biosynthesis, and TAG content was increased up to 32.5-fold in 8-week-old leaf blades. However, the ectopic expression of *BdWRI1* also caused cell death in leaves, which has not been previously observed in dicotyledonous plants such as *Arabidopsis* (*Arabidopsis thaliana*) (Yang et al., 2015). Lipid analysis indicated that the free FA content was 2.0-fold elevated in *BdWRI1*-expressing leaf blades of *Brachypodium*. The transcription of predicted genes involved in  $\beta$ -oxidation was induced. In addition, linoleic fatty acid treatment caused cell death in *Brachypodium* leaf blades, an effect

that was reversed by addition of the FA biosynthesis inhibitor cerulenin. Taken together, ectopic expression of *BdWR11* in *Brachypodium* enhances FA biosynthesis and TAG accumulation in leaves as expected but also leads to increased free FA content, which has cytotoxic effects leading to cell death. Thus, while *WR11* appears to ubiquitously affect FA biosynthesis and TAG accumulation in diverse plants, its ectopic production can lead to undesired side effects depending on the context of the specific lipid metabolism of the respective plant species.

This work was funded in part by the US Department of Energy Great Lakes Bioenergy Research Center Cooperative Agreement DE-FC02-07ER64494 and MSU AgBioResearch.

## References

- Cernac A, Andre C, Hoffmann-Benning S, Benning C (2006) *WR11* is required for seed germination and seedling establishment. *Plant Physiol* 141: 745-757
- Cernac A, Benning C (2004) *WRINKLED1* encodes an AP2/EREB domain protein involved in the control of storage compound biosynthesis in *Arabidopsis*. *Plant J* 40: 575-585
- Ma W, Kong Q, Arondel V, Kilaru A, Bates PD, Thrower NA, Benning C, Ohlrogge JB (2013) *Wrinkled1*, a ubiquitous regulator in oil accumulating tissues from *Arabidopsis* embryos to oil palm mesocarp. *PLoS One* 8: e68887
- Ma W, Kong Q, Grix M, Mantyla JJ, Yang Y, Benning C, Ohlrogge JB (2015) Deletion of a C-terminal intrinsically disordered region of *WRINKLED1* affects its stability and enhances oil accumulation in *Arabidopsis*. *Plant J* 83: 864-874
- Sanjaya, Durrett TP, Weise SE, Benning C (2011) Increasing the energy density of vegetative tissues by diverting carbon from starch to oil biosynthesis in transgenic *Arabidopsis*. *Plant Biotechnol J* 9: 874-883
- Sanjaya, Miller R, Durrett TP, Kosma DK, Lydic TA, Muthan B, Koo AJ, Bukhman YV, Reid GE, Howe GA, Ohlrogge J, Benning C (2013) Altered lipid composition and enhanced nutritional value of *Arabidopsis* leaves following introduction of an algal diacylglycerol acyltransferase 2. *Plant Cell* 25: 677-693
- Yang Y, Munz J, Cass C, Zienkiewicz A, Kong Q, Ma W, Sanjaya, Sedbrook J, Benning C (2015) Ectopic Expression of *WRINKLED1* Affects Fatty Acid Homeostasis in *Brachypodium distachyon* Vegetative Tissues. *Plant Physiol* 169: 1836-1847