

The Secretome of *Caldicellulosiruptor bescii*: Biomass Deconstruction without Conventional Pretreatment

Jenna Young^{1,4*} (jmy715@uga.edu), Daehwan Chung,^{1,4} Joseph Groom,^{1,4} Minsoek Cha,^{1,4} Debra Mohnen,^{1,4} Richard Dixon,^{2,4} Yannick J. Bomble,^{3,4} Michael E. Himmel,^{3,4} Janet Westpheling,^{1,4} and **Paul Gilna**⁴

¹University of Georgia, Athens; ²University of North Texas, Denton; ³National Renewable Energy Laboratory, Golden, Colorado; ⁴BioEnergy Science Center, Oak Ridge National Laboratory, Oak Ridge, Tennessee

<http://bioenergycenter.org>

Project Goals: The BioEnergy Science Center (BESC) is focused on the fundamental understanding and elimination of biomass recalcitrance. BESC's approach to improve accessibility to the sugars within biomass involves (1) designing plant cell walls for rapid deconstruction and (2) developing multi-talented microbes or converting plant biomass into biofuels in a single step [consolidated bioprocessing (CBP)]. BESC research in biomass deconstruction and conversion targets CBP by studying thermophilic anaerobes to understand novel strategies and enzyme complexes for biomass deconstruction and manipulating these microorganisms for improved conversion, yields, and biofuel titer.

Members of the bacterial genus *Caldicellulosiruptor* are the most thermophilic cellulolytic organisms so far described and have the ability to grow on lignocellulosic biomass without conventional pretreatment. A comparison of the pangenome of this genus suggests that among the genes important for cellulolytic ability are *celA* and a cluster of genes involved in pectin degradation (1). CelA is a bi-functional glycoside hydrolase that contains a Family 9 endoglucanase and a Family 48 exoglucanase joined by three Family 3 carbohydrate binding modules (CBMs), and while there are two Family 9 and three Family 48 glycoside hydrolases in *C. bescii*, CelA is the only protein that combines both activities. Deletion of CelA resulted in a significant decrease in cellulolytic activity (2). Expression of full length CelA in *C. bescii* revealed that extracellular CelA protein is glycosylated whereas intracellular CelA is not (3). The mechanism and role of protein glycosylation in bacteria is poorly understood and the ability to express CelA *in vivo* in *C. bescii* will allow the study of the mechanism of protein glycosylation in this thermophile. Moreover, heterologous expression of an additional thermophilic endoglucanase (E1) from *Acidothermus cellulolyticus*, enhanced the ability of *C. bescii* to deconstruct biomass demonstrating that while extremely effective, the ability of *C. bescii* to deconstruct plant biomass can be improved (4).

The *C. bescii* genome contains five genes predicted to be involved in pectin deconstruction/ utilization and three exist in a cluster with a predicted transcriptional regulator. Expression of this cluster is significantly up-regulated in cells growing on biomass (5). Most biomass models do not list pectin because of its low abundance in grass walls and in dicot secondary walls (6). Recent work has shown that pectin is synthesized in secondary walls (7), that some pectin biosynthetic enzymes are amplified in grasses (8), and that saccharification of plant biomass can be improved by modifying the structure of pectin (9). Deletion of the pectinase gene cluster in *C. bescii* resulted in a mutant reduced in its ability to grow on dicot and grass biomass (10). The phenotype of the *C. bescii* pectinase mutant provides direct genetic evidence that pectin is a significant barrier to

deconstruction of untreated plant biomass by *C. bescii* and that pectin plays an important and underappreciated role in biomass recalcitrance.

References

1. Blumer-Schuetz SE, Giannone RJ, Zurawski JV, Ozdemir I, Ma Q, Yin YB, et al., "*Caldicellulosiruptor* Core and Pangenomes Reveal Determinants for Noncellulosomal Thermophilic Deconstruction of Plant Biomass," *Journal of Bacteriology* 2012;194(15):4015-28.
2. Young J, Chung D, Bomble YJ, Himmel M, Westpheling J., "Deletion of *Caldicellulosiruptor bescii* CelA Reveals its Crucial Role in the Deconstruction of Lignocellulosic Biomass," *Biotechnology for Biofuels*. 2014;7:142.
3. Chung D, Young J, Bomble YJ, Vander Wall TA, Groom J, Himmel ME, et al., "Homologous Expression of the *Caldicellulosiruptor bescii* CelA Reveals that the Extracellular Protein Is Glycosylated," *PloS One* 2015;10(3):e0119508. Epub 2015/03/24.
4. Chung D, Young J, Cha M, Brunecky R, Bomble YJ, Himmel ME, et al., "Expression of the *Acidothermus Cellulolyticus* E1 Endoglucanase in *Caldicellulosiruptor bescii* Enhances its Ability to Deconstruct Crystalline Cellulose," *Biotechnology for Biofuels* 2015;8:113. Epub 2015/08/14.
5. Kataeva I, Foston MB, Yang SJ, Pattathil S, Biswal AK, Poole FL, et al., "Carbohydrate and Lignin are Simultaneously Solubilized from Untreated Switchgrass by Microbial Action at High Temperature," *Energy and Environmental Science* 2013;6(7):2186-95.
6. Mohnen D, Bar-Peled M, Somerville C., "Cell Wall Polysaccharide Synthesis," *Biomass Recalcitrance*: Blackwell Publishing Ltd.; 2009. p. 94-187.
7. Atmodjo MA, Sakuragi Y, Zhu X, Burrell AJ, Mohanty SS, Atwood JA, III, et al., "Galacturonosyltransferase (GAUT)1 and GAUT7 are the Core of a Plant Cell Wall Pectin Biosynthetic Homogalacturonan:Galacturonosyltransferase Complex," *Proceedings of the National Academy of Sciences of the United States of America* 2011;108(50):20225-30. Epub 2011/12/03.
8. Yin Y, Chen H, Hahn MG, Mohnen D, Xu Y., "Evolution and Function of the Plant Cell Wall Synthesis-Related Glycosyltransferase Family 8," *Plant Physiology* 2010;153(4):1729-46. Epub 2010/06/05.
9. Lionetti V, Francocci F, Ferrari S, Volpi C, Bellincampi D, Galletti R, et al., "Engineering the Cell Wall by Reducing De-Methyl-Esterified Homogalacturonan Improves Saccharification of Plant Tissues for Bioconversion," *Proceedings of the National Academy of Sciences of the United States of America* 2010;107(2):616-21. Epub 2010/01/19.
10. Chung D, Pattathil S, Biswal AK, Hahn MG, Mohnen D, Westpheling J., "Deletion of a Gene Cluster Encoding Pectin Degrading Enzymes in *Caldicellulosiruptor bescii* Reveals an Important Role for Pectin in Plant Biomass Recalcitrance," *Biotechnology for Biofuels* 2014;7(1):147. Epub 2014/10/18.

The BioEnergy Science Center is a U.S. Department of Energy Bioenergy Research Center supported by the Office of Biological and Environmental Research in the DOE Office of Science.