

## Exploring Species Specificity of Lambda Red Recombination

Gabriel Filsinger<sup>1\*</sup> ([filsinger@g.harvard.edu](mailto:filsinger@g.harvard.edu)), Tim Wannier,<sup>1</sup> Xavier Rios,<sup>1</sup> Chris Gregg,<sup>1</sup> Marc Lajoie,<sup>2</sup> **George M. Church<sup>1</sup>**

<sup>1</sup>Harvard Medical School, Department of Genetics, Boston, MA; <sup>2</sup>University of Washington, Seattle, WA

<http://arep.med.harvard.edu>

**Project Goals: To extend recombineering methods to organisms other than *E. coli* by identifying and overcoming the host-specificity of Beta recombinases.**

Recombination using the lambda red protein Beta has been extensively used in *E. coli* to incorporate insertions, deletions, and point mutations into the genome at chosen loci. Although Beta increases the rate of ssDNA recombination in *E. coli* 10,000 fold, this catalytic recombination activity has not been observed in other bacteria such as *Lactobacillus* or *Corynebacterium*. Interestingly, the family of single-stranded annealing proteins that includes Beta has been found to have species-specific activities and unpredictable efficiencies, making it difficult to design a generalized method for porting Beta recombination across organisms. In order to explore mechanisms of generalizing the recombinase activity, we use *Lactobacillus* and *E. coli* as two model organisms with orthogonal recombination systems to probe the source of species specificity.

*This project has been funded by DOE grant DE-FG02-02ER63445 and a NSF Graduate Research Fellowship.*