

## High-Resolution Spatial Analysis Reveals How Nitrogen Source Governs Carbon Partitioning Between Members in a Phototrophic Consortium

Christopher R. Anderton<sup>1\*</sup> (christopher.anderton@pnnl.gov), Ryan S. Renslow,<sup>1</sup> Jennifer M. Mobberley,<sup>1</sup> William B. Chrisler,<sup>1</sup> Jessica K. Cole,<sup>1</sup> Amy A. Boaro,<sup>1</sup> Jamie Nuñez,<sup>1</sup> Yasemin Yesiltepe,<sup>1</sup> Beau Morton,<sup>1</sup> Alexandra Cory,<sup>1</sup> Hayley Cardamone,<sup>1</sup> Mary S. Lipton,<sup>1</sup> James Moran,<sup>1</sup> Stephen R. Lindemann,<sup>2</sup> Jim Fredrickson,<sup>1</sup> and **Lee Ann McCue**<sup>1</sup>

<sup>1</sup>Pacific Northwest National Laboratory, Richland, WA; <sup>2</sup>Purdue University, West Lafayette, IN

<http://www.pnnl.gov/biology/programs/fsfa/>

**Project Goals: The goal of the Metabolic and Spatial Interactions in Communities (MOSAIC) Foundational Scientific Focus Area is to understand the fundamental mechanisms by which microbial metabolic interactions and spatial organization impact carbon, nitrogen, and energy dynamics in microbial communities. Our studies focus on the coupling of carbon and nitrogen cycles in microbial communities, the role of environmental variables in governing the rates of these cycles, and the impact of environmental perturbations on microbial community dynamics. We employ tractable model consortia whose member genome sequences have been defined, advanced omics measurements, functional imaging, taxonomic profiling, and modeling to elucidate interaction mechanisms within complex microbial communities. Our research supports the DOE goals to achieve a predictive understanding of Earth's integrated biogeochemical processes.**

Interactions between microbial photoautotrophs and associated heterotrophic organisms are ubiquitous in nature and exert significant impacts on global biogeochemical cycling. Hence, elucidating the molecular mechanisms by which environmental conditions impact community interactions in phototrophic-heterotrophic consortia is critical to predicting how they will respond to environmental change.<sup>1,2</sup> Here, we used a model consortium containing one photoautotrophic cyanobacterium (*Phormidium* sp. OSCR) and 18 associated heterotrophic species, for which a species-resolved metagenome reconstruction is available,<sup>3</sup> to understand fundamental carbon and nitrogen coupling in microbial communities. We hypothesized that the genome-predicted inability of most heterotrophic consortium members to directly acquire  $\text{NO}_3^-$  from the medium would result in widespread nitrogen limitation for these species. Altering the available nitrogen source ( $\text{NO}_3^-$  vs.  $\text{NO}_3^- + \text{NH}_4^+$ ) drastically altered community composition and dynamics, though not in ways easily predictable from members' ability to assimilate  $\text{NO}_3^-$ . Examination of the consortia, using a multimodal mass spectrometry-based stable isotope probing approach, revealed that although  $\text{NH}_4^+$  was acquired approximately twice as rapidly as  $\text{NO}_3^-$ , there was no significant difference in community  $\text{H}^{13}\text{CO}_3^-$  incorporation. To resolve whether C/N partitioning among members was altered, we used high-lateral resolution mass spectrometry imaging (NanoSIMS), in conjunction with our previously developed image processing pipeline<sup>4</sup>. These data showed significant differences in the rate at which heterotrophs

acquired cyanobacterially-fixed carbon, where carbon was much more rapidly transferred to heterotrophs when  $\text{NO}_3^-$  was the sole nitrogen source than when  $\text{NH}_4^+$  was also available. Relating these results to that from proteomic analysis of the cyanobacteria indicates altered iron acquisition and pyruvate metabolism based on the available nitrogen source, suggesting that carbon transfer rate may be increased when  $\text{NO}_3^-$  is the sole nitrogen source as the cyanobacteria are discarding excess reductant. Fluorescence *in situ* hybridization (FISH) imaging with species-specific probes provided insight into the potential role of select heterotrophs within the community. We observed that the most abundant heterotroph (*Aliidiomarina calidilacus*), independent of nitrogen source, was always proximal to the cyanobacterial filaments, whereas the heterotroph (*Algoriphagus marincola*) whose abundance was significantly reduced upon  $\text{NH}_4^+$  amendment co-localized with lysed cyanobacterial cells. These results suggest that *A. calidilacus* cells form an epibiotic or parasitic relationship with the cyanobacteria, and that *A. marincola* cells are detritivores that recycle proteinaceous biomass.

## References

- (1) Cole, J.; Hutchison, J.; Renslow, R.; Kim, Y.-M.; Chrisler, W.; Engelmann, H.; Dohnalkova, A.; Hu, D.; Metz, T.; Fredrickson, J.; Lindemann, S. *Frontiers in Microbiology* **2014**, *5*.
- (2) Lindemann, S.; Moran, J.; Stegen, J.; Renslow, R.; Hutchison, J.; Cole, J.; Dohnalkova, A.; Tremblay, J.; Singh, K.; Malfatti, S.; Chen, F.; Tringe, S.; Beyenal, H.; Fredrickson, J. *Frontiers in Microbiology* **2013**, *4*.
- (3) Nelson, W. C.; Maezato, Y.; Wu, Y.-W.; Romine, M. F.; Lindemann, S. R. *Applied and Environmental Microbiology* **2016**, *82*, 255-267.
- (4) Renslow, R. S.; Lindemann, S. R.; Cole, J. K.; Zhu, Z.; Anderton, C. R. *Biointerphases* **2016**, *11*, 02A322.

*This research was supported by the U.S. Department of Energy (DOE), Office of Biological and Environmental Research (BER), as part of BER's Genomic Science Program (GSP). This contribution originates from the GSP Foundational Scientific Focus Area (FSFA) at the Pacific Northwest National Laboratory (PNNL). A portion of this work was performed in the William R. Wiley Environmental Molecular Sciences Laboratory (EMSL), a national scientific user facility sponsored by OBER and located at PNNL. PNNL is a multi-program national laboratory operated by Battelle for the DOE under Contract DE-AC05-76RLO 1830. The work conducted by the U.S. Department of Energy Joint Genome Institute (JGI) was supported by the Office of Science of DOE under Contract No. DE-AC02-05CH11231 and Community Sequencing Project 701.*