

***Rhodospiridium toruloides* for conversion of depolymerized cellulose, hemicellulose, and lignin into bioproducts**

Junko Yaegashi^{1,2*} (JYaegashi@lbl.gov), James Kirby^{1,3}, Masakazu Ito⁴, Jian Sun^{1,5}, Tanmoy Dutta^{1,5}, Mona Mirsiaghi⁶, Eric R. Sundstrom⁶, Edward Baidoo^{1,6}, Deepti Tanjore⁶, Todd Pray⁶, Kenneth Sale^{1,5}, Seema Singh^{1,5}, Jay D. Keasling^{1,3,6,7,8}, Blake A. Simmons^{1,6}, Steven W. Singer^{1,6}, Adam P. Arkin^{4,6,8}, Jeffrey M. Skerker^{4,6}, **Jon K. Magnuson^{1,2}**, **John M. Gladden^{1,5}**

¹ Joint BioEnergy Institute, Emeryville, CA, USA; ² Pacific Northwest National Laboratory, WA, USA; ³ California Institute of Quantitative Biosciences (QB3), University of California, Berkeley, CA, USA; ⁴ Energy Bioscience Institute, Berkeley, CA, USA; ⁵ Sandia National Laboratory, Livermore, CA, USA; ⁶ Lawrence Berkeley National Laboratory, Berkeley, CA, USA; ⁷ Department of Chemical & Biomolecular Engineering, University of California, Berkeley, CA, USA; ⁸ Department of Bioengineering, University of California, Berkeley, CA, USA.

<https://www.jbei.org/research/divisions/deconstruction/fungal-biotechnology/>

Project Goals: Conversion of all major components of lignocellulose into advanced bioproducts has been a long elusive goal that is essential for enabling a robust bioeconomy. This project, performed in the Fungal Biotechnology group in the Deconstruction Division at JBEI, seeks to establish and utilize *Rhodospiridium toruloides* as a platform organism that can convert depolymerization products of cellulose, hemicellulose and lignin into biofuels and bioproducts.

Economical conversion of lignocellulosic biomass to biofuels and bioproducts is central to the establishment of a robust bioeconomy. Efficient conversion of lignocellulose requires the utilization of all its primary components (cellulose, hemicellulose, and lignin), yet no microbe in commercial use today can achieve this feat. To that end, we explored the utility of *Rhodospiridium toruloides* as a new platform organism for the production of biofuels and bioproducts from deconstructed plant biomass. In this study, *Rhodospiridium toruloides* was engineered to produce two non-native terpenes with biofuel (bisabolene) and pharmaceutical (amorphadiene) applications from a mixture of depolymerized cellulose (glucose), hemicellulose (xylose), and lignin (*p*-coumaric acid). The titers of these bioproducts in synthetic defined media with 2% glucose were 294mg/L and 36g/L, respectively. It was also cultivated on corn stover hydrolysates prepared by two different pretreatment methods, including one using the novel biocompatible ionic liquid choline α -ketoglutarate. This organism was able to tolerate the ionic liquid that was used for pretreatment and was also able to utilize all of the three major carbon sources in the hydrolysate to support growth and bisabolene production. Finally, we demonstrate that this organism is amenable to high-gravity fed-batch fermentation, reaching a titer of 680mg/L in alkaline corn stover hydrolysate. This study establishes *R. toruloides* as a new platform for the simultaneous conversion of depolymerized cellulose, hemicellulose, and lignin into biofuels and bioproducts.

This work was performed as part of the DOE Joint BioEnergy Institute (<http://www.jbei.org>) supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research, through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U.S. Department of Energy.