

Characterization of Plant Golgi-Localized Nucleotide Sugar Transporters

Julien Sechet^{1,2}, Beibei Jing^{1,2}, Soe Htwe^{1,2}, Fekadu Anderbehan^{1,2}, Noriko Inada³, **Jenny C Mortimer**^{1,2*} (jcmortimer@lbl.gov)

¹Joint BioEnergy Institute, Emeryville, CA; ²Biosciences Area, Lawrence Berkeley National Laboratory, Berkeley, CA; ³The University of Tokyo, Japan

<http://jbei.org/>

Project Goals: The most abundant organic material on earth is lignocellulosic biomass or non-food plant material. JBEI's mission is to convert biomass to biofuels. The goal is to provide the nation with clean, renewable transportation fuels identical to gasoline, diesel and jet fuel. Building a successful lignocellulosic biofuels industry depends, in part, on developing specialized biofuel crops or feedstocks that are optimized for deconstruction into sugars and fermentation into biofuels and bioproducts.

Abstract: The majority of the nucleotide sugar substrates required for plant polysaccharide biosynthesis are made in the cytosol. Non-cellulosic polysaccharides, including hemicelluloses and pectins, are synthesized inside the Golgi lumen by glycosyltransferases. Nucleotide sugar transporters (NSTs) are therefore required to move the substrates to the correct compartment. Whilst Arabidopsis is estimated to have ~50 NSTs, only 4 are predicted to transport GDP-sugars. Arabidopsis synthesizes four different GDP-sugars: GDP-D-mannose, GDP-D-glucose, GDP-L-galactose and GDP-L-fucose which glycosylate pectins, hemicelluloses, proteins and sphingolipids.

Recent work has shown that the Arabidopsis GOLGI NUCLEOTIDE SUGAR TRANSPORTER1 (GONST1) specifically transports GDP-D-mannose for sphingolipid glycosylation¹ and GOLGI FUCOSE TRANSPORTER (GFT) transports GDP-L-fucose for multiple purposes, including xyloglucan fucosylation². Here, we have characterized the remaining two predicted GDP-sugar transporters: GONST2 and GONST3, and identified specific functions for both. These data will be discussed in the context of engineering plant cell walls for improved production of biofuels and biochemicals.

References

1. Mortimer et al. (2013) Plant Cell 25: 1881.
2. Rautengarten et al. (2016) Nature Communications 7:12119.

Funding statement.

This work was part of the DOE Joint BioEnergy Institute (<http://www.jbei.org>) supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research,

through contract DE-AC02-05CH11231 between Lawrence Berkeley National Laboratory and the U. S. Department of Energy