

A Concerted Systems Biology Analysis of Aromatic Metabolism in

***Rhodococcus opacus* PD630**

Garrett W. Roell^{1*}, Rhiannon R. Carr¹, Tayte Campbell², Zeyu Shang¹, William R. Henson¹, Jeffrey Czajka¹, **Hector Garcia Martin**^{3,4,5,6}, **Fuzhong Zhang**¹, **Marcus Foston**¹, **Gautam Dantas**^{2,7,8,9}, **Tae Seok Moon**¹, and **Yinjie J. Tang**¹

¹Department of Energy, Environmental and Chemical Engineering, Washington University in St. Louis, St. Louis, MO, 63130, USA; ²The Edison Family Center for Genome Sciences and Systems Biology, Washington University in St. Louis School of Medicine, St. Louis, MO, 63110, USA; ³DOE, Joint BioEnergy Institute, Emeryville, CA, 94608, USA; ⁴DOE, Agile BioFoundry, Emeryville, CA, 94608, USA; ⁵Biological Systems and Engineering Division, Lawrence Berkeley National Lab, Berkeley, CA, 94720, USA; ⁶BCAM, Basque Center for Applied Mathematics, Bilbao, Spain; ⁷Department of Pathology and Immunology, Washington University in St. Louis School of Medicine, St. Louis, MO, 63108, USA; ⁸Department of Biomedical Engineering, Washington University in St. Louis, St Louis, MO, 63130, USA; ⁹Department of Molecular Microbiology, Washington University in St. Louis School of Medicine, St. Louis, MO, 63108, USA

Project Goals:

1. Use ^{13}C -MFA to reveal *R. opacus*' phenol metabolism
2. Connect flux data with transcription profiling and metabolite analysis to show phenol metabolism regulations.
3. Determine whether *R. opacus* phenol utilization is hindered by other aromatic and non-aromatic substrates.
4. Test adaptively evolved strains to determine how their central metabolic network has changed.

Rhodococcus opacus PD630 metabolizes aromatic substrates and naturally produces branched-chain lipids, which are advantageous traits for lignin valorization. To provide insights into its lignocellulose hydrolysate utilization, we performed ^{13}C pathway tracing, transcriptional profiling, biomass composition analysis, and metabolite profiling in conjunction with ^{13}C -metabolic flux analysis (MFA) of phenol metabolism. We found that 1) phenol is metabolized through the ortho branch of the β -ketoacid pathway; 2) phenol-fed cultures have high TCA cycle fluxes with overflow succinate secretion; 3) NADPH is generated mainly via NADPH-dependent isocitrate dehydrogenase; 4) Active cataplerotic fluxes increase plasticity in the TCA cycle; and 5) gluconeogenesis occurs partially through the reversed Entner–Doudoroff pathway (EDP). We also found that phenol-fed *R. opacus* PD630 generally has lower sugar phosphate concentrations (e.g., fructose 1,6-bisphosphatase < 5%) compared to metabolite pools in glucose-fed *Escherichia coli* (set as 100%), while pool sizes of its TCA metabolites (malate, succinate, and α -ketoglutarate) are higher than those in *E. coli*. In addition, glucose catabolite repression is absent in *R. opacus*, but phenol utilization can be hindered by the presence of other aromatic substrates (e.g., benzoate). Three adaptively-evolved strains display different growth rates when fed with phenol as a sole carbon source, but they demonstrate a conserved central flux network.

This project is supported by the DoE BER grant (DESC0018324).